

AD-AU929/2- BECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)	
REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
I. REPORT NUMBER 2, GOV ACCESSION NO). RECIPIENT'S CATALOG NUMBER
Technical Report, No. 2, / Ang 77-20	NOU EUS
Induction of Olefin Metathesis by Acetylenes.	Technical Report, 8/1/79- 11/20/80
	6. PERFORMING ORG. REPORT NUMBER
Thomas J./Katz, Steven J./Lee/Mridula/Nair and/Edward B./Savage	8. CONTRACT OR GRANT NUMBER(*) NOQ014-79-C-9683
PERFORMING ORGANIZATION NAME AND ADDRESS Thomas J. Katz	16. PROGRAM ELEMENT, PROJECT, TASI AREA & WORK UNIT NUMBERS
Chemistry Department Columbia University, New York, N.Y. 10027	NR 356-726
1. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
Office of Naval Research - Chemistry Arlington, VA 22217	November 1989
4. MONITORING AGENCY NAME & ADRESS AGENCY Gentrelling Office)	Unclassified
LEVELY LEVELY	154. DECLASSIFICATION/DOWNGRADING
This document has been approved for public release. 7. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different for	
To be published in the <u>Journal of the American Ch</u>	emical Society
. KEY WORDS (Continue on reverse side if necessary and identify by block number	
Polymers Metathesis Alkenes Polymerization Metal-carbenes Alkynes Catalysis Polyacetylenes Initiation Polyalkenamers	
ABSTRACT (Continuo en reverso side il necessary and identify by block number) Acetylenes induce (phenylmethoxycarbene)pentacarbe metatheses of cis-2-pentene, cyclooctene, cyclohe too much quenches the reactions. The first three remarkably stereoselective, yielding olefins that	onyltungsten to initiate ptene, and cyclopentene, but of these transformations are

observations accord with the hypothesis that Scheme 1 is the mechanism of acetylene polymerization. They also provide a functional equivalent (an acetylene) of the organometallic cocatalysts like ethylaluminum dichloride, unique in operating in the absence of strong Lewis acid metal-halides.

DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

4//337 8012 08 03

OFFICE OF NAVAL RESEARCH

Contract NOO014-79-C-0683

Task No. NR 356-726

TECHNICAL REPORT NO. 2

Induction of Olefin Metathesis by Acetylenes

bу

Thomas J. Katz, Steven J. Lee, Mridula Nair and Edward B. Savage

Prepared for Publication

in the

Journal of The American Chemical Society

Columbia University
Department of Chemistry
New York, New York

November 20, 1980

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited

Induction of Olefin Metathesis by Acetylenes

Sir:

If the reason that metal-carbenes polymerize acetylenes is because the transformations involve the steps in Scheme 1.1,2 then combining metal-

Scheme 1

carbenes and acetylenes should generate other, possibly more reactive metalcarbenes that might initiate transformations their precursors could not. We recently discovered that acetylenes can induce olefin metatheses. 3,4 and we are reporting here how the process can be made to work generally, and remarkably stereospecifically. We have discovered that acetylenes not only induce the reactions, but quench them as well, and in an accompanying paper we analyze why. 5

Table 1 shows that if 200 equivalents of three cycloalkenes are combined with 1 equivalent of (phenylmethoxycarbene)pentacarbonyltungsten 1,0 no polyalkenamers 1a,7 form, but that if phenylacetylene is present, the reactions (eq 1) do indeed take place, even if slowly. However, preliminary attempts to effect the analogous transformation with the seemingly related molecule cis-2-pentene were ineffective, even when ever larger amounts of acetylenes and large amounts of metal-carbenes were used,

Avallability

Accession

NATS

$$c = w(co)_5$$
 $c = w(co)_5$
 $c = w(co)_5$
 $c = w(co)_5$

as initiators. Thus 200 equivalents of phenylacetylene and 7 of 1 after 6

days at 50 °C converted 160 equivalents of cis-2-pentene into less than half an equivalent of the butenes plus hexenes although 67% of the phenylace-tylene was transformed and isolated as its polymer.

The key to bringing about the reactions more efficiently, it turns out, is to use less (not more) of the acetylene. Fig. 1 illustrates this for combinations of cyclopentene (100 equiv.), phenylacetylene (varying amounts), and metal-carbene 1 (1 equivalent), for it shows that the inverse of the yield of polypentenamer increases linearly with the concentration of phenylacetylene. Notice according to this graph, that to achieve a 100% yield, one simply has to use no phenylacetylene at all!

Similar increases in yields with decreasing acetylene concentration have been measured for other olefins and are illustrated by the enhanced rates of conversion (compared to the ones described above) in Table 2.

<u>Table 1</u>. Yields of Polyalkenamers obtained in Metatheses Induced by Combining Cycloalkenes (200 equiv.), Phenylacetylene (10 equiv.), and Metal-Carbenes 1 (1 equiv.) at 50 °C. $^{\frac{a}{2}}$

Cycloalkene b	Time (days)	Yield ^c ,d (%)	
5	7	19	
7	13	16	
8	24	7	

In the absence of phenylacetylene, the three cycloalkenes after 7, 26, and 24 days respectively gave no polyalkenamers. b Number of carbons. C The yields are those of the polyalkenamers isolated by dissolving the reaction products in CH₂Cl₂, precipitating with CH₃OH, and drying in a vacuum. They may be in error by ca. 20% (relative). d After 5 days neither cycloheptene nor cyclooctene polymerized appreciably (< ca. 6%), for the reaction mixtures were not yet significantly viscous.

Table 2. Yields and Molecular Weights of Products Obtained in Metatheses Induced by Combining Olefins (100 equiv.), Phenylacetylene (1 equiv.), and Metal-Carbene 1 (1 equiv.) at 50 °C.

Olefin	Time (days)	Yield (%)	™ = x10 ⁻³	ਔ ≜ n-3
cycloheptene	2	16 <u>b</u>	321	135
cyclooctene	3.5	8 <u>p</u>	281	147
cis-2-pentene	7	7.9⊆		

Molecular weights were analyzed by gel permeation chromatography in toluene on Waters Associates μ -styragel. The values recorded are half the weights of the polystyrenes that would exhibit the chromatograms observed (see references 5 and 9). $\frac{b}{2} \pm \frac{ca}{2}$ 20% (relative). $\frac{c}{2}$ 100 (moles butene + moles hexene)/ moles pentene. The error is $\frac{ca}{2}$. 4% (relative).

These metatheses can be very highly stereoselective. ¹⁰ For cycloheptene (IR analysis: ¹¹ 97±1% cis, ¹³CNMR: ¹² >96%), ¹⁴ cyclooctene (IR: 94.3±2% cis, ¹³CNMR: 94±1%), ¹⁵ and cis-2-pentene (the 2-butenes are 96.5± 0.6% cis, the 3-hexenes 93.7±0.8% cis), ¹⁸ although not for cyclopentene (IR: 74±8% cis, ¹³CMR: 58%), ²⁰ the stereoselectivities are substantially higher than for every other initiator previously studied, with one significant exception, (diphenylcarbene)pentacarbonyltungsten (2). ⁹, ¹⁹ This is a remarkable observation, yet it agrees with the introductory premises because the effect of the transformation in Scheme 1 is to convert metal-carbene ½ into a substance whose structure is essentially that of 2. The acetylene is thus an activator, but unlike the organometallic cocatalysts like C₂H₅AlCl₂, which it replaces, it is unique in operating in the absence of strong Lewis acid metal-halides, whose presence may account for why most metatheses are only weakly stereoselective ¹⁹, ²¹ and for why some are accompanied by undesirable side-reactions. ²²


The observation that acetylenes induce metatheses supports the idea that acetylene polymerization like olefin metathesis involves the addition of metal-carbenes to carbon-carbon multiple bonds. The observation that acetylenes quench metatheses too is also in accord with the hypothesis as analyzed in the following paper. 5

Acknowledgment. We are grateful for the support of the National Science Foundation (CHE-77-22726) and the U.S. Office of Naval Research.

Thomas J. Katz. Steven J. Lee,
Mridula Nair. Edward B. Savage
Department of Chemistry,
Columbia University
New York, New York 10027

Figure Caption

Figure 1. Inverse of the yield of polypentenamer as a function of the ratio of phenylacetylene and cyclopentene. Cyclopentene (100 equiv.), phenylacetylene (varying amounts), and 1 (1 equiv.) were combined at 50 °C for 21.5 h. The bars show the errors introduced if the weight of the polymer obtained from 680 mg of cyclopentene is measured incorrectly by 10 mg. The yield (100%) at [phenylacetylene] = 0 is derived from theory (see reference 5). The slope and standard deviation are those of the least squares straight line through the first four points, weighted as the inverses of the relative errors in the inverse yields. The last three points were discarded because when the yields and molecular weights are low, the fraction of polymer lost in the isolation can be high. (These points therefore all appear above the line.)



References and Notes

- (1) Katz, T.J.; Lee, S.J. J. Am. Chem. Soc., 1980, 102, 422.
- (2) Masuda, T.; Sasaki, N.; Higashimura, T. Macromolecules 1975, 8, 717.
- (3) (a) Katz, T.J. Adv. Organomet. Chem. 1977, 16, 283. (b) Grubbs, R.H. Prog. Inorg. Chem. 1978, 24, 1. (c) Rooney, J.J.; Stewart, A. Catalysis (London) 1977, 1, 277. (d) Calderon, N.; Lawrence, J.P.; Ofstead, E.A. Adv. Organomet. Chem. 1979, 17, 449. (e) Haines, R.J.; Leigh, G.J. Chem. Soc. Rev. 1975, 4, 1. (f) Mol, J.C.; Moulijn, J.A. Adv. Catal. 1975, 24, 131.
- (4) Some of the results were discussed at the 3rd NSF Workshop on Organo-metallic Chemistry, Pingree Park, Colorado, July 16, 1979, and at the 3rd International Symposium on Olefin Metathesis, Lyons, France, September 12, 1979. See Katz, T.J.; Lee, S.J.; Shippey, M.A. J. Molec. Catal. 1980, 8, 219.
- (5) Katz, T.J.; Savage, E.B.; Lee, S.J.; Nair, M. J. Am. Chem. Soc., 1980, accompanying manuscript.
- (6) (a) Fischer, E.O.; Maasböl, A. Chem. Ber. 1967, 100, 2445. (b) Aumann, R.; Fischer, E.O. Angew. Chem. Int. Ed. Engl. 1967, 6, 878. (c) Darensbourg, D.J. Inorg. Chem. 1970, 9, 32.
- (7) Dall'Asta, G. Rubber Chem. Technol. 1974, 47, 511.
- (8) The form of this relationship is discussed in reference 5.
- (9) Katz, T.J.; Lee, S.J.; Acton, N. Tetrahedron Lett. 1976, 4247.
- (10) One set of samples was prepared by combining 100 equiv. of each olefin, 1 equiv. acetylene, and 1 equiv. 1 at 50 °C. Other samples were prepared using somewhat different ratios, and for cyclooctene and cycloheptene some were prepared at 65 °C. No obvious difference was noticed in the measurements of their stereochemistries, which are averaged here.

- (11) Measured by the ratio of the intensities of the infrared peaks at \underline{ca} . 1400 and 960 cm⁻¹. See reference 9, footnote 16.
- (12) The intensities of the trans and cis C-2 resonances (see reference 9, the references in its footnote 18, and reference 13) were measured, and for polyoctenamer those of the two olefinic carbon resonances as well.
- (13) (a) Carmen, C.J.; Wilkes, C.E. <u>Macromolecules</u> <u>1974</u>, <u>7</u>, 40. (b) Chen, H.Y. <u>Appl. Polym. Spectrosc.</u> <u>1978</u>, <u>7</u>.
- (14) Except in reference 9, polyheptenamer has previously been obtained 6-39% cis in 3-18% yield: (a) Natta, G.; Dall'Asta, G.; Bassi, I.;

 Carella, G. Makromol. Chem. 1966, 91, 87. (b) Porri, L.; Diversi, P.;

 Lucherini, A.; Rossi, R. ibid. 1975, 176, 3121. (c) Brit. Pat. 1 062 367, 1967.
- (15) WCl₆ (or in one case a tungsten-carbene) + organoaluminum compounds have given polyoctenamers that are largely trans. 14a,b,16 At low temperatures in the presence of certain modifiers [BF₃·(C₂H₅)₂O at -30 °C or dissobutylaluminum oxide at +12 °C] or when WF₆ is used in place of WCl₆, the polymer formed is said to be largely cis. 17
- (16) (a) Calderon, N.; Morris, M. J. Polym. Sci. Part A-2 1967, 5, 1283.
 (b) Höcker, H.; Musch, R. Makromol. Chem. 1974, 175, 1395. (c) Arlie,
 J.-P.; Chauvin, Y.; Commercuc, D.; Soufflet, J.-P. 1bid. 1974, 175, 861.
- (17) (a) Syatkowsky, A.I.; Denisova, T.T.; Ikonitsky, I.V.; Babitsky, B.D.
 J. Polym. Sci., Polym. Chem. Ed. 1979, 17, 3939. (b) Zimmerman, M.;
 Lehnert, G.; Maertens, D.; Pampus, G. Ger. Offen. 2 334 604, 1975.
 (c) Dall'Asta, G.; Manetti, R. Ital. Pat. 932 461, 1971, mentioned in reference 7, p.551. (d) Castner, K.F. U.S. Pat. 4 038 471, 1977; Chem.
 Abstr. 1978, 87, 152726a.

- (18) The stereochemistries were measured for the reactions of 2-pentene when the extents of metathesis were between 6 and 8%. Except in reference 19, 2-butene has never been obtained more than 83% cis (and that only at "zero time"; more commonly it is 58% cis at zero time). The 3-hexene is formed even less stereospecifically. See the references in footnote 2 of reference 19. Also (a) Leconte, M.; Basset, J.M.

 J. Am. Chem. Soc. 1979, 101, 7296. (b) Leconte, M.; Ben Taarit, Y.;
 Bilhou, J.L.; Basset, J.M. J. Molec. Catal. 1980, 8, 263.
- (19) Katz, T.J.; Hersh, W.H. Tetrahedron Lett. 1977, 585.
- (20) The polypentamer formed by 2 is, as here, not as highly cis as the other polyalkenamers. 9
- (21) For additional indications that the stereoselectivity of cyclopentene's metathesis falls with added aluminum compounds and with temperature see (a) K.J. Ivin, D.T. Laverty, and J.J. Rooney, Makromol. Chem., 178, 1545 (1977) and (b) I.A. Oreshkin, Bull. Acad. Sci. USSR, Div. Chem. Sci., 26, 2377 (1977); Eur. Polym. J., 13, 447 (1977).
- (22) (a) Hocks, L.; Noels, A.; Hubert, A.; Teyssie, P. J. Org. Chem. 1976, 41, 1631 and references therein. (b) Wolovsky, R.; Moaz, N.; Nir, Z. Synthesis 1970, 2, 656. (c) Graham, J.R.; Slaugh, L.H. Tetrahedron Lett. 1971, 787. (d) Zowade, T.; Höcker, H. Makromol. Chem. 1973, 165, 31. (e) Wang, J.L.; Menapace, H.R. J. Org. Chem. 1968, 33, 3794; (f) Menapace, H.R.; Maly, N.A.; Wang, J.L.; Wideman, L.G. J. Org. Chem. 1975, 40, 2983. (g) Vialle, J.; Basset, J.M. Reaction Kinetics and Catal. Lett. 1975, 2, 397.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research		U.S. Army Research Office	
Attn: Code 472		Attn: CRD-AA-IP	
800 North Quincy Street		P.O. Box 1211	
Arlington, Virginia 22217	2	Research Triangle Park, N.C. 27709	1
ONR Branch Office		Naval Ocean Systems Center	
Attn: Dr. George Sandoz		Attn: Mr. Joe McCartney	•
536 S. Clark Street	•	San Diego, California 92152	1
Chicago, Illinois 60605	1	Novel Massage Contag	
ONR Area Office		Naval Weapons Center Attn: Dr. A. B. Amster,	
		Chemistry Division	
Attn: Scientific Dept.		China Lake, California 93555	1
715 Broadway New York, New York 10003	1	China bake, Calliothia 93333	•
New lolk, New lolk 10005	•	Naval Civil Engineering Laboratory	
ONR Western Regional Office		Attn: Dr. R. W. Drisko	
1030 East Green Street		Port Hueneme, California 93401	1
Pasadena, California 91106	1	,,,,,,,	-
rabatina, ourrename vero	-	Department of Physics & Chemistry	
ONR Eastern/Central Regional Office		Naval Postgraduate School	
Attn: Dr. L. H. Peebles		Monterey, California 93940	1
Building 114, Section D		• •	
666 Summer Street		Dr. A. L. Slafkosky	
Boston, Massachusetts 02210	1	Scientific Advisor	
		Commandant of the Marine Corps	
Director, Naval Research Laboratory		(Code RD-1)	
Attn: Code 6100		Washington, D.C. 20380	1
Washington, D.C. 20390	1		
		Office of Naval Research	
The Assistant Secretary		Attn: Dr. Richard S. Hiller	
of the Navy (RE&S)		800 N. Quincy Street	_
Department of the Navy		Arlington, Virginia 22217	1
Room 4E736, Pentagon	_		
Washington, D.C. 20350	1	Naval Ship Research and Development Center	
Commander, Naval Air Systems Command	l	Attn: Dr. G. Bosmajian, Applied	
Attn: Code 310C (H. Rosenwasser)		Chemistry Division	
Department of the Navy		Annapolis, Maryland 21401	1
Washington, D.C. 20360	1		
		Naval Ocean Systems Center	
Defense Technical Information Center		Attn: Dr. S. Yamamoto, Marine	
Building 5, Cameron Station		Sciences Division	,
Alexandria, Virginia 22314	12	San Diego, California 91232	1
Dr. Fred Saalfeld		Mr. John Boyle	
Chemistry Division, Code 6100		Materials Branch	
Naval Research Laboratory		Naval Ship Engineering Center	_
Washington, D.C. 20375	1	Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, 356B

	No. Copies		No. Copies
Dr. T. C. Williams		Douglas Aircraft Company	
Union Carbide Corporation		Attn: Technical Library	
Chemical and Plastics		Cl 290/36-84	
Tarrytown Technical Center		AUTO-Sutton	
Tarrytown, New York	1	3855 Lakewood Boulevard	
•		Long Beach, California 90846	1
Dr. R. Soulen			
Contract Research Department		NASA-Lewis Research Center	
Pennwalt Corporation		Attn: Dr. T. T. Serafini, MS 49-1	
900 First Avenue		21000 Brookpark Road	_
King of Prussia, Pennsylvania 19406	1	Cleveland, Ohio 44135	1
Dr. A. G. MacDiarmid		Dr. J. Griffith	
University of Pennsylvania		Naval Research Laboratory	
Department of Chemistry		Chemistry Section, Code 6120	_
Philadelphia, Pennsylvania 19174	1	Washington, D.C. 20375	1
Dr. C. Pittman		Dr. G. Goodman	
University of Alabama		Globe-Union Incorporated	
Department of Chemistry		5757 North Green Bay Avenue	
University, Alabama 35486	1	Milwaukee, Wisconsin 53201	1
Dr. H. Allcock		Dr. E. Fischer, Code 2853	
Pennsylvania State University		Naval Ship Research and	
Department of Chemistry		Development Center	
University Park, Pennsylvania 16802	1	Annapolis Division	
		Annapolis, Maryland 21402	1
Dr. M. Kenney			
Case-Western University		Dr. Martin H. Kaufman, Head	_
Department of Chemistry		Materials Research Branch (Code 4542))
Cleveland, Ohio 44106	1	Naval Weapons Center	_
		China Lake, California 93555	1
Dr. R. Lenz			
University of Massachusetts		Dr. J. Magill	
Department of Chemistry	_	University of Pittsburg	
Amherst, Massachusetts 01002	1	Metallurgical and Materials	
		Engineering	•
Dr. M. David Curtis		Pittsburg, Pennsylvania 22230	1
University of Michigan		D. 0 411	
Department of Chemistry	•	Dr. C. Allen	
Ann Arbor, Michigan 48105	1	University of Vermont	
Dm. M. Cood		Department of Chemistry	1
Dr. M. Good		Burlington, Vermont 05401	•
Division of Engineering Research		Dr. D. Rozobreitor	
Louisiana State University	1	Dr. D. Bergbreiter Texas A&M University	
Baton Rouge, Louisiana 70803	1	Department of Chemistry	
		College Station, Texas 77843	1
			-

TECHNICAL REPORT DISTRIBUTION LIST, 356B

No.	
Copie	s

Professor R. Drago Department of Chemistry University of Illinois Urbana, Illinois 61801

1

Dr. F. Brinkman
Chemical Stability & Corrosion
Division
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

,

Professor H. A. Titus Department of Electrical Engineering Naval Postgraduate School Monterey, California 93940

1

COL R. W. Bowles, Code 100M Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217

Arlington, Virginia 22217

Professor T. Katz
Department of Chemistry
Columbia University
New York, New York 10027

1

Professor James Chien
Department of Chemistry
University of Massachusetts
Amherst, Massachusetts 01002

Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta, Georgia 30314

1

SP472-3/A3

472:GAN:716:ddc 78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No. Copies

Dr. Rudolph J. Marcus Office of Naval Research Scientific Liaison Group American Embassy APO San Francisco 96503

1

Mr. James Kelley DTNSRDC Code 2803 Annapolis, Maryland 21402

1